



ecSatter is Correlated with Superior Wellness Indicators

Compiled by Ellyn Satter March 14, 2022

At the heart of a good relationship with food is the principle of Eating Competence.¹ Eating Competence, as defined by the Satter Eating Competence Model (ecSatter), is being positive, comfortable, and flexible with eating as well as matter-of-fact and reliable about getting enough to eat of personally enjoyable, nourishing food. Even though ecSatter says nothing at all about what or how much to eat or what to weigh, people who are Eating Competent (EC) have better diets, lower BMIs, superior metabolic profiles, more-positive quality of life indicators, are more active, do better with respect to managing a limited food budget, and do better with respect to parenting with food. Eating competence is measured by the validated ecSI 2.0^{2,3} across all income levels.⁴

EC showed superior diets

Even though ecSatter doesn't emphasize eating "healthy" food and avoiding "unhealthy" food, and, instead, gives strong permission to eat preferred food, people who are EC do better nutritionally and with FV availability.

- Pandemic-related EC decrease (**31.69 vs 29.99** $p < 0.005$) correlated with decreased FV and increased sugary beverage consumption.⁵
- EC parents of fourth graders showed superior dietary quality (HEI scores) and a flexible definition of mealtime regularity.⁶
- EC predominantly non-Hispanic parents of fourth graders had more in-home FV, modeled more healthful eating, higher FV efficacy ("I can prepare fruit that my child will eat"), higher modeling ("How often do you prepare food at home?"), and had more in-home FV availability.⁷
- EC Finnish adults had better diet quality (<0.001).⁸
- EC Finnish adults at risk of diabetes ate more fruits and vegetables.⁹
- EC Finnish adolescents showed greater family meal frequency, more frequent consumption of FV, and more health-promoting family eating patterns.⁹
- EC Hispanic parents tended to have food-preparation skills and positive attitudes toward cooking, cook with their child, eat breakfasts and dinners with their child, include and enjoy vegetables at dinnertime, and have greater in-home FV availability.¹⁰
- Low-income EC had higher fiber, vitamin A, E, C, most B-vitamins, magnesium, iron, zinc, potassium, higher Healthy Eating Index, were more likely to follow the Prudent dietary pattern.¹¹
- EC low-income were most likely to plan meals, include all the food groups, shop with a list, cook from scratch, offer and serve more fruit.²
- EC university students had highest perceived dietary quality.¹²
- EC Spanish elderly ate more fruit, were more adherent to Mediterranean diet.¹³
- EC mixed income were most likely to offer and serve more FV, and were in action maintenance stage of change with FV.³

EC showed superior wellness, medical nutrition therapy measures

Even though ecSatter gives strong permission to eat preferred food and encourages using fat, salt, and sugar to make food taste good, people who are EC show superior wellness and metabolic indicators.

- EC had lower previously undiagnosed type 2 diabetes, metabolic syndrome, HbA1c, triglycerides ($p < 0.05$ for all), lower BMI and fasting insulin, higher HDL (<0.001).⁸
- Low-income EC had fewer oral health issues.¹⁴
- EC Spanish elderly had higher HDL-cholesterol, lower fasting blood glucose, total cholesterol, LDL/HDL cholesterol, diabetes, family HX early-onset heart disease.¹³
- EC mixed-income adults had higher HDL-cholesterol, lower systolic and diastolic blood pressure, even when stress tested (speech and cold-water immersion) and lower LDL, triglycerides.¹⁵

EC showed lower BMI

Even though ecSatter does not encourage striving in any way for weight loss and, instead, encourages eating as much as is desired of preferred food and letting weight find its own level, people who are EC have the same or lower BMIs. *Starred (*)*: *Weight taken by trained personnel*.

- Pandemic-related EC decrease EC (**31.69 vs 29.99** $p < 0.005$) correlated with weight increase.⁵
- EC parents of fourth graders have lower BMI.⁶
- *EC Finnish adults at risk for diabetes have lower BMI, are less likely to be obese.⁸
- EC Hispanic parents of fourth graders had lower incidence of overweight, obesity.⁶
- EC Finnish adolescents more often perceived their weight as appropriate and less often had tried to lose weight.⁹
- *EC university students have lowest incidence of overweight/obesity.^{16, 17}
- *EC university students have lowest BMI.¹²
- EC low-income female adults have lower BMI.²
- *EC elderly Spanish have lower BMI.¹³
- EC subjects in validation trial have lower BMI.³

EC did better with low-income food management

Even though ecSatter offers no particular guidance on budgeting or food resource management, low-income EC tend to see themselves as food secure, whereas equally low-income non-EC are more likely to characterize themselves as food insecure. EC do better with respect to managing limited food resources.

- Low-income EC don't self-identify as food insecure. ecSI 2.0 scores: High/marginal food security: 30.4; low food security: 23.1. ¹⁴
- EC least likely to run out of food before end of month, most likely to feel confident about managing food money.²
- Low-income EC least likely to self-identify as food insecure, most likely to plan meals, shop with a list, cook from scratch.³

EC showed superior quality of life indicators

Even though ecSatter is about eating, EC has to do with quality of life. EC captures positive eating attitudes and behaviors, which both reflect and act as a proxy for other quality of life indicators. EC, mental health, and self-care all depend on detecting, respecting, and responding to feelings and preferences. Direct measures of quality of life (sleep, activity), mental health (emotional, psychological, and social well-being), and physical self-esteem all correlate with EC.

- EC parents scored high on sDOR.2-6y and had higher parent quality of life indicators (sleep, stress, emotional, psychological, and social well-being).¹⁸
- EC Finnish adults at risk of diabetes had higher physical activity and better sleep quality, were less likely to smoke.⁸
- EC Finnish adolescents more often perceived their body size as appropriate, had less often tried to lose weight, had higher self-esteem and a stronger sense of coherence.⁹
- EC university students showed more physical activity, higher sleep duration¹⁶ and quality of sleep.¹⁷
- EC introductory nutrition students were less likely to report past or current eating disorders.¹⁹
- EC low-income adult females showed lowest interpersonal distrust, impulsivity, ineffectiveness, maturity fears and social insecurity as well as lowest body dissatisfaction, drive for thinness, restrained eating, disinhibited eating and (intolerable) hunger.^{2,3}
- EC university students had lowest desired weight loss, lowest emotional eating, lowest psychological/ emotional distress.¹²
- EC university students had higher weight satisfaction, lower desire to lose weight.²⁰
- EC low-income adults in cognitive interviews had positive eating attitudes and behaviors: food seeking, enjoyment, satisfaction, relaxation, positive re cooking. **Non-EC** low-income had negative eating attitudes and behaviors: emphasis on weight management, food restriction and avoidance, sporadic and/or inattentive eating, guilt/misery regarding eating the wrong food, negativity regarding cooking, desire to buy low-fat and diet products, desire to stop eating junk food.²¹

EC showed higher activity

Even though ecSatter says nothing about activity, EC have higher levels of physical activity and indicators of activity.

- EC Finnish adults had higher physical activity.⁸
- EC college students showed correlations across EC tertiles for >60 minutes of physical activity.¹⁶
- EC low-income women showed correlations across EC tertiles for self-reported physical activity.²
- EC college students reported higher levels of physical active, had a higher VO₂ max, a measure of lung function.¹²
- EC adults across income categories showed correlations across EC tertiles for self-reported physical activity.³

EC showed better parenting with food

Even though ecSatter says nothing about feeding children, EC parents do better with respect to following the Satter Division of Responsibility in Feeding (sDOR) and raising children who have lower nutritional risk.

- Video-captured mothers of 2 through 5-year-olds who indicated on sDOR.2-6y that they followed sDOR indicated they actually did take leadership and give autonomy with feeding.²²
- EC parents scored higher on sDOR.2-6y and have children with lower nutritional risk, apply less restriction and pressure on child's eating.¹⁸
- EC showed highest fdSI (precursor to sDOR.2-6y), are likely to divide feeding responsibilities, view child's eating positively.²³
- EC mothers of 2 through 5-year-olds showed low restriction, appropriately divided feeding responsibilities.²⁴

References

1. Satter E. Eating Competence: definition and evidence for the Satter Eating Competence Model. *J Nutr Educ Behav*. 2007;39:S142-S153.
2. Krall JS, Lohse B. Validation of a measure of the Satter Eating Competence model with low-income females. *Int J Behav Nutr Phys Act*. 2011;8. doi:10.1186/1479-5868-8-26 PMC3094263,
3. Lohse B, Satter E, Horacek T, et al. Measuring Eating Competence: psychometric properties and validity of the ecSatter Inventory. *J Nutr Educ Behav*. 2007;39:S154-S166.
4. Godleski S, Lohse B, Krall JS. Satter Eating Competence Inventory subscale restructure after confirmatory factor analysis. *J Nutr Educ Behav*. Jul 23 2019;
5. Queiroz FL, Nakano EY, Botelho RB, et al. Eating Competence among Brazilian Adults: A comparison between before and during the COVID-19 Pandemic. *Foods*. 2021;10. doi:10.3390/foods10092001
6. Lohse B, Faulring K, Mitchell DC, et al. A definition of "regular meals" driven by dietary quality supports a pragmatic schedule. *Nutrients*. 2020;12. doi:10.3390/nu12092667
7. Lohse B, Pflugh Prescott M, Cunningham-Sabo L. Eating Competent parents of 4th grade youth from a predominantly non-Hispanic white sample demonstrate more healthful eating behaviors than non-eating competent parents. *Nutrients*. 2019;11. doi:10.3390/nu11071501 PMC6682872,
8. Tilles-Tirkkonen T, Aittola K, Männikkö R, et al. Eating Competence is associated with lower prevalence of obesity and better insulin sensitivity in Finnish adults with increased risk for type 2 diabetes: The StopDia Study. *Nutrients*. 2019;12. doi:10.3390/nu12010104 PMC7019577,
9. Tilles-Tirkkonen T, Nuutinen O, Suominen S, et al. Preliminary Finnish measures of Eating Competence suggest association with health-promoting eating patterns and related psychobehavioral factors in 10–17 year old adolescents. *Nutrients*. 2015;7:3828-3846.
10. Lohse B, Cunningham-Sabo L. Eating Competence of Hispanic parents is associated with attitudes and behaviors that may mediate fruit and vegetable-related behaviors of 4th grade youth. *J Nutr* 2012;142:1903-1909.
11. Lohse B, Bailey RL, Krall JS, et al. Diet quality is related to Eating Competence in cross-sectional sample of low-income females surveyed in Pennsylvania. *Appetite*. 2012;58:645-650.
12. Greene GW, Schembre SM, White AA, et al. Identifying clusters of college students at elevated health risk based on eating and exercise behaviors and psychosocial determinants of body weight. *J Am Diet Assoc*. 2011;111:394-400

References (Cont'd)

13. Lohse B, Psota T, Estruch R, et al. Eating Competence of elderly Spanish adults is associated with a healthy diet and a favorable cardiovascular disease risk profile. *J Nutr*. 2010;140:1322-1327.
14. Lohse B, Masters L. Eating Competence and oral health in Supplemental Nutrition Assistance Program eligible populations. *J Dent Hyg*. 2019;93:42-50.
15. Psota TL, Lohse B, West SG. Associations between Eating Competence and cardiovascular disease biomarkers. *J Nutr Educ Behav*. 2007;39:S171-S178.
16. Quick V, Shoff S, Lohse B, et al. Relationships of eating competence, sleep behaviors and quality, and overweight status among college students. *Eat Behav*. 2015;19:15-19.
17. Quick V, Byrd-Bredbenner C, White A, et al. Eat, sleep, work, play: associations of weight status and health-related behaviors among young adult college students. *Am J Health Promot*. 2013:e64-e72. doi:10.4278/ajhp.130327-QUAN-130.
18. Lohse B, Mitchell DC. Valid and reliable measure of adherence to Satter Division of Responsibility in Feeding. *J Nutr Educ Behav*. 2021:211-222.
19. Brown LB, Larsen KJ, Nyland NK, et al. Eating competence of college students in an introductory nutrition course. *J Nutr Educ Behav*. 2013;45:269-273.
20. Clifford D, Linda A, Keeler LA, et al. Weight attitudes predict Eating Competence among college students. *Fam Consum Sci Res J*. 2010;39:184-193.
21. Stotts Krall J, Lohse B. Interviews with low-income Pennsylvanians verify a need to enhance eating competence. *J Am Diet Assoc*. 2009;109:468-473.
22. Lohse B, Satter E. Use of an observational comparative strategy demonstrated construct validity of a measure to assess adherence to the Satter Division of Responsibility in Feeding. *J Acad Nutr Diet*. 2021;121:1143-1156.e6.
23. Lohse B, Satter E, Arnold K. Development of a tool to assess adherence to a model of the division of responsibility in feeding young children: using response mapping to capacitate validation measures. *Child Obes*. 2014;10:153-168.
24. Tylka TL, Eneli IU, Kroon Van Diest AM, et al. Which adaptive maternal eating behaviors predict child feeding practices? An examination with mothers of 2- to 5-year-old children. *Eat Behav*. 2013;14:57-63.